



# Spirulina platensis shields the submandibular gland from cadmium toxicity by bolstering antioxidant defenses and maintaining its structural integrity

Eman Mohamed Faruk<sup>a,b,\*</sup>, Fatma Ibrahim<sup>c</sup>, Dalia Ibrahim El-wafaey<sup>d</sup>,  
Yasmeen Mohammed Ismail El Sayed<sup>e</sup>, Mohamed Ghazy Attia Hablas<sup>f</sup>, Mahmoud M. Hassan<sup>g</sup>,  
Ahmed Mohammed Zaazaa<sup>h</sup>, Kamal M. Kamal<sup>i</sup>

<sup>a</sup> Anatomy Department, College of Medicine, Umm Al-Qura University, Makkah, Saudi Arabia

<sup>b</sup> Department of Histology and Cell Biology, Faculty of Medicine, Benha University, Benha, Egypt

<sup>c</sup> Department of Forensic Medicine and Clinical Toxicology, Faculty of Medicine, Zagazig University, Zagazig, Egypt

<sup>d</sup> Department of Human Anatomy and Embryology, Faculty of Medicine, Zagazig University, Egypt

<sup>e</sup> Department of Pharmacology, Faculty of Medicine, Benha University, Benha, Egypt

<sup>f</sup> Department of Histology and Cell Biology, Faculty of Medicine, Suez University, Egypt

<sup>g</sup> Department of Physiology, Faculty of Medicine, Benha University, Benha, Egypt

<sup>h</sup> Student at Faculty of Medicine, Benha National University, Benha Colleges in Cairo, Egypt

<sup>i</sup> Department of Anatomy and Embryology, Faculty of Medicine, Benha University, Benha, Egypt

## ARTICLE INFO

Editor: Elisabetta Liverani

### Keywords:

Cadmium chloride  
*Spirulina platensis*  
Submandibular gland  
Interleukin 1-beta

## ABSTRACT

Cadmium (Cd), an element categorized as a non-essential transitional metal, has potential hazards to the health of both human beings and animals. *Spirulina platensis* (SP), a type of blue-green algae, possesses a high concentration of essential antioxidants. The present study aimed to explore the possible defensive role of SP against Cd-induced submandibular gland injury in rats by assessment of biomarkers related to both oxidative stress and inflammatory processes, which were further explored through histopathological examination of submandibular gland tissue. Consequently, the study included 32 mature rats, subdivided into four different groups as follows: control, SP, Cadmium chloride (CdCl<sub>2</sub>), and CdCl<sub>2</sub>/SP. The duration of the study was 24 days. The results revealed that CdCl<sub>2</sub> induced submandibular gland injury as shown by the oxidant/antioxidant imbalance and increased inflammatory reactions, in addition to, histopathological changes and overexpression of BAX immunostaining. Concurrent SP administration to CdCl<sub>2</sub>-treated rats significantly improved all these effects. We concluded that concurrent SP supplement improved the submandibular gland injury provoked by CdCl<sub>2</sub>.

## 1. Introduction

Cadmium (Cd), an element categorized as a non-essential transitional metal, possesses toxic properties and potential hazards to the health of both human beings and animals as well, it is noteworthy to mention that the toxicity of Cd remains remarkably high, even at extremely minute concentrations [1]. It is a naturally occurring environmental pollutant that is produced by agricultural and manufacturing sources [2]. Cd is largely used in several industries such as paints, metal coating, mining, batteries, and fertilizers [3].

Cadmium is toxic both by inhalation and ingestion causing acute and chronic toxicities. Some edible foodstuffs exhibit selective uptake of Cd, therefore food is frequently described as a primary route through which humans are exposed to Cd [4]. Food and tobacco smoking together are

the most significant sources of Cd for human beings [5].

Cd accumulates in nearly all of the human tissues with a biological half-life that has been assessed to be more than 10 years [6]. It was evidenced that consuming water from natural sources leads to the high Cd accumulation in the renal cortex and the liver of individuals living in the industrial regions [7]. Cd in blood is an indicator of recent exposure, while Cd in urine mirrors the body burden and reveals chronic exposure [8].

The generation of reactive oxygen species (ROS) for instance superoxide anion, hydrogen peroxide, and hydroxyl radical, induced by Cd, results in the activation of signaling pathways. This, in turn, stimulates autophagy, apoptosis, and gene expression, denoting that oxidative stress is a chief mechanism of Cd toxicity [9]. Also, Cd targets the sulfhydryl groups present in proteins causing its inactivation, which can

\* Corresponding author at: Faculty of Medicine, Department of Histology and Cell Biology, Benha University, 44519, Egypt.

E-mail address: [faruk\\_eman@yahoo.com](mailto:faruk_eman@yahoo.com) (E.M. Faruk).

result in multiple functional defects in the nuclei, the endoplasmic reticulum, and mitochondria [10]. Cd is a mitogen and promotes cancer in several tissues. It is proven to be a human carcinogen (group I of the International Agency for Research on Cancer classification) [11].

The equilibrium between oxidant formation and endogenous antioxidant protective mechanisms is required. If this balance is disturbed, it can lead to oxidative stress-induced tissue damage [12].

*Spirulina platensis* (SP), a type of blue-green algae, possesses a high concentration of essential antioxidants, for instance, phycocyanin, carotenoids, phenolic compounds, microelements, vitamins, amino acids, and fatty acids besides antioxidant enzymes, such as superoxide dismutase, catalase, and peroxidase [13,14].

*Spirulina platensis* usage is well-established in prophylaxis besides treatment of several toxicity models, for example, acrylamide [15], fluoride [16], cyclophosphamide [17], arsenic [18], lead [19], mercury [20], manganese [21], and cadmium [22].

Considering this perspective, this study aimed to recognize the defensive properties of SP against CdCl<sub>2</sub>-triggered submandibular gland impairment. As far as our knowledge extends, this is the foremost study that has been conducted to explore these effects.

## 2. Material and methods

### 2.1. Tested compounds and chemicals

Cadmium chloride (CdCl<sub>2</sub>) was acquired from Sigma–Aldrich Chemicals Co., St. Louis, USA (Catalogue number: 202908). *Spirulina platensis* (SP), in the form of dark greenish fine particles, was obtained from El-hellow for biological products, Cairo, Egypt. Normal saline (0.9 % sodium chloride), used as a vehicle, was obtained from Nile Company, Cairo.

### 2.2. Animals and study design

The study encompassed 32 adult male albino rats, with an age range of 6–8 weeks and weight range of 180–200 gm, obtained from the Zagazig Scientific and Medical Research Center (ZSMRC), Faculty of Medicine, Zagazig University, where the study had been conducted also. The present study has been approved by the Institutional Animal Care and Use Committee (IACUC) at Zagazig University, as evidenced by the assigned approval number: ZU-IACUC/3/F/413/2022. All experimental techniques were executed in agreement with the guidelines established for the ethical treatment and use of laboratory animals.

After two weeks of acclimatization to guarantee their physical health and to exclude any sick animals, the rats were weighed and divided into equal four groups, with eight rats in each.

Group I (Control): rats were given normal saline (1 ml) that was used also as a vehicle to dissolve SP and CdCl<sub>2</sub>.

Group II (SP): rats were given 300 mg/kg/b.w. of SP [23].

Group III (CdCl<sub>2</sub>): rats were given 10 mg/kg/b.w. of CdCl<sub>2</sub> [24,25].

Group IV (CdCl<sub>2</sub>/SP): rats were given SP (300 mg/kg/b.w.) and CdCl<sub>2</sub> (10 mg/kg/b.w.).

All treatments were given by oral gavage, once/day for successive 24 days.

### 2.3. Blood and submandibular tissue preparation

Upon completion of the treatment protocol, the body weight of every rat was documented to accurately assess any weight changes.

Blood samples were collected using clean test tubes and then centrifuged for 5 min for separation of the serum. The supernatant sera were used to measure malondialdehyde (MDA), catalase (CAT), superoxide dismutase (SOD), reduced glutathione (GSH), and total antioxidant capacity (TAC). Then, all rats were scarified via cervical dislocation, and the submandibular glands were dissected and weighed. The submandibular glands were separated into two parts. The first one

was homogenized for measurement of transforming growth factor-beta (TGF-β), tumor necrosis factor-alpha (TNF-α), Interleukin 1-beta (IL-1β), and Interleukin-10 (IL-10). The other part was prepared for histological studies.

### 2.4. Determination of the activity of oxidant and antioxidant biomarkers

The levels of MDA (CAT. No. MD 25 29), CAT (CAT. No. CA 25 17), SOD (CAT. No. SD 25 21), GSH (CAT. No. GR 25 11) and TAC (CAT. No. TA 25 13) were measured by the specific kits purchased from Biodiagnostic, Egypt.

### 2.5. Determination of inflammatory markers in submandibular tissue

Based on enzyme-linked immuno-sorbent assay (ELISA), the levels of TGF-β, TNF-α, IL-1β, and IL-10 were assessed by the commercial ELISA kits purchased from (eBioscience, Inc., San Diego, CA, USA) as stated by manufacturer's guidelines.

### 2.6. Histological study

#### 2.6.1. Light microscopy

The submandibular glands tissues were dissected and fixed in 10 % formalin for processing of paraffin sections. 5 μm-thickness sections that were prepared and processed for H&E staining and PAS staining (for detection of mucopolysaccharides) [26] then examined by a light microscope. We performed comprehensive histological analyses for regional differences in the acini, myoepithelium, ducts, blood vessels, and the extracellular matrix (ECM) surrounding acini and ducts. Serous acini, showing abundant basophilic cytoplasm with a round nucleus, and mucous acini with slightly basophilic cytoplasm.

#### 2.6.2. Immunohistochemistry

For the immunohistochemistry, the method of peroxidase-labelled streptavidin–biotin was employed [27]. It was done to detect the Bax proteins. The antibodies employed in the detection of Bax consisted of rabbit polyclonal antibodies obtained from Sigma-Aldrich, Egypt, and were utilized at a 1:50 dilution rate. The specific Catalog No. for the antibodies was SAB4502549. The reaction was shown as brownish cytoplasmic discoloration. For negative control sections, the primary antibodies were excluded. Positive control of BAX was taken from a human breast cancer specimen.

#### 2.6.3. Histomorphometric analysis

The morphometric analysis of the sections was conducted utilizing the Leica Qwin 500 Image Analyzer Computer System, located at the Pathology Department of the Faculty of Medicine at Cairo University. To evaluate the mean area percentage of PAS staining and Bax immunostaining among the studied groups, ten distinct fields were evaluated in each slide, at lower magnification. The histological parameters were semi-quantitatively defined by essential histological features, including the degree of fatty replacement, preservation of lobular architecture, preservation of ducts and acini, the presence of interstitial fibrosis, and the inflammatory component. The scales used were no degeneration (0), trace (1), mild (2), moderate (3), and severe (4) degeneration.

The non-parametric Cuzick's test for trend [28] was used to compare the grades of fatty replacement, loss of lobular architecture, presence of interstitial fibrosis, degree of ductal degeneration, presence of diffuse inflammatory component, presence of focal inflammatory component, and degree of acinar degeneration across these four groups.

Each parameter was scored (1–4 points), so the total point score for each rat was ranging between 6 and 24. The obtained scores were divided by six for interpretation. Less than 1: no degeneration, 1.0–2.5: mild degeneration, and 2.6–4.0: severe degeneration.

## 2.7. Statistical analysis

The variables were presented in the form of mean  $\pm$  standard deviation (SD) and were subjected to comparison through the application of one-way analysis of variance (ANOVA) and post-Hoc Tukey test to determine the differences between groups. Differences were considered significant when the *P*-value was less than 0.05. IBM SPSS software was used for statistical analysis.

## 3. Results

### 3.1. SP effectively attenuated CdCl<sub>2</sub> toxicity affecting mortality rates, body weight, and the weight of the submandibular gland

There were no recorded fatalities within any of the studied groups. CdCl<sub>2</sub> treatment for successive 24 days resulted in a statistically significant body weight loss compared with the other studied groups (*P* < 0.05). Regarding the submandibular gland weight, there was no significant difference in the CdCl<sub>2</sub>/SP when compared to the control and SP groups (*P* > .05) (Table 1).

### 3.2. SP remarkably restored CdCl<sub>2</sub>-induced oxidative stress and inflammatory biomarkers imbalance in the rats

This study revealed that there were no statistically significant differences in mean values of all studied biochemical parameters in the SP group when compared with the control group (*P* > .05).

CdCl<sub>2</sub>-intoxicated rats showed increased oxidative stress represented by statistically significantly higher MDA levels and lower catalase, SOD, GSH, and TAC activities compared with the other studied groups (*P* < 0.05). Concurrent SP administration to CdCl<sub>2</sub>-intoxicated rats induced a significant decrease in the MDA level and an increase in catalase, SOD, GSH, and TAC activities (Table 2).

The results of Table 3 showed that CdCl<sub>2</sub> significantly increased TGF- $\beta$ , TNF- $\alpha$ , and IL-1 $\beta$  levels along with significantly decreased IL-10 levels. The SP administration had protected against these alterations represented by a significant lowering of the levels of TGF- $\beta$ , TNF- $\alpha$ , and IL-1 $\beta$  and a significant elevation of IL-10 level.

### 3.3. SP alleviated CdCl<sub>2</sub>-induced submandibular histopathological and histochemical alternation in the rats

Histological analysis of H&E-stained sections of the submandibular salivary gland in the control and SP groups showed normal parenchyma with many lobules that contain both pale vacuolated mucous acini with flat nuclei and serous acini (with rounded nuclei and darkly stained secretory granules) and serous demilunes (capped acini). Ducts were found between acini in which intralobular, and the intercalated ducts were characterized by their simple cuboidal epithelium lining, while the striated ducts were distinguished by their lining of high cuboidal cells.

**Table 1**

Effects of cadmium chloride and *Spirulina platensis* administrations on the initial body weight, final body weight, and submandibular gland weight.

Parameter	Groups			
	Control	SP	CdCl <sub>2</sub>	CdCl <sub>2</sub> /SP
Initial body weight (gm)	185.38 $\pm$ 5.10	187.38 $\pm$ 3.29	185.50 $\pm$ 3.67	184.63 $\pm$ 2.67
Final body weight (gm)	239.13 $\pm$ 3.64	241.38 $\pm$ 6.48	203.25 $\pm$ 4.89 <sup>ab</sup>	224.75 $\pm$ 3.06 <sup>abc</sup>
Submandibular weight (gm)	0.43 $\pm$ 0.02	0.42 $\pm$ 0.01	0.35 $\pm$ 0.03 <sup>ab</sup>	0.40 $\pm$ 0.04 <sup>c</sup>

Each value is mean  $\pm$  SD for eight rats in each group.

<sup>a</sup>: *P* < 0.05 compared to control, <sup>b</sup>: *P* < 0.05 compared to SP, <sup>c</sup>: *P* < 0.05 compared to CdCl<sub>2</sub>.

Abbreviations; SP, *Spirulina platensis*; CdCl<sub>2</sub>, Cadmium chloride.

**Table 2**

Effects of cadmium chloride and *Spirulina platensis* administrations on the oxidant and antioxidant biomarkers.

Parameter	Groups			
	Control	SP	CdCl <sub>2</sub>	CdCl <sub>2</sub> /SP
Serum MDA (nmol/ml)	1.55 $\pm$ 0.30	1.52 $\pm$ 0.32	3.93 $\pm$ 0.41 <sup>ab</sup>	2.97 $\pm$ 0.12 <sup>abc</sup>
Serum Catalase (U/L)	129.03 $\pm$ 2.20	127.62 $\pm$ 3.56	66.91 $\pm$ 4.11 <sup>ab</sup>	99.29 $\pm$ 3.19 <sup>abc</sup>
Serum SOD (U/ml)	371.73 $\pm$ 9.29	368.19 $\pm$ 3.52	95.70 $\pm$ 6.72 <sup>ab</sup>	176.04 $\pm$ 6.80 <sup>abc</sup>
Serum GSH (mmol/L)	5.08 $\pm$ 0.12	5.02 $\pm$ 0.15	1.48 $\pm$ 0.18 <sup>ab</sup>	3.52 $\pm$ 0.18 <sup>abc</sup>
Serum TAC (mmol/L)	1.88 $\pm$ 0.12	1.84 $\pm$ 0.11	0.64 $\pm$ 0.04 <sup>ab</sup>	0.93 $\pm$ 0.06 <sup>abc</sup>

Each value is mean  $\pm$  SD for eight rats in each group.

<sup>a</sup>: *P* < 0.05 compared to control, <sup>b</sup>: *P* < 0.05 compared to SP, <sup>c</sup>: *P* < 0.05 compared to CdCl<sub>2</sub>.

Abbreviations; MDA, Malondialdehyde; SOD, Superoxide dismutase; GSH, Reduced glutathione; TAC, Total antioxidant capacity; SP, *Spirulina platensis*; CdCl<sub>2</sub>, Cadmium chloride.

**Table 3**

Effects of cadmium chloride and *Spirulina platensis* administrations on the pro-inflammatory and fibrotic markers.

Parameter	Groups			
	Control	SP	CdCl <sub>2</sub>	CdCl <sub>2</sub> /SP
Tissue TGF- $\beta$ (pg/100 mg tissue)	175.75 $\pm$ 4.81	177.43 $\pm$ 1.63	412.87 $\pm$ 5.24 <sup>ab</sup>	330.88 $\pm$ 5.27 <sup>abc</sup>
Tissue TNF- $\alpha$ (pg/100 mg tissue)	20.94 $\pm$ 1.26	21.13 $\pm$ 1.82	54.51 $\pm$ 4.61 <sup>ab</sup>	37.75 $\pm$ 2.93 <sup>abc</sup>
Tissue IL-10 (pg/100 mg tissue)	377.22 $\pm$ 6.63	372.03 $\pm$ 5.24	133.01 $\pm$ 4.60 <sup>ab</sup>	234.22 $\pm$ 4.62 <sup>abc</sup>
Tissue IL-1 $\beta$ (pg/100 mg tissue)	86.54 $\pm$ 1.06	88.08 $\pm$ 1.09	314.02 $\pm$ 2.54 <sup>ab</sup>	250.43 $\pm$ 5.01 <sup>abc</sup>

Each value is mean  $\pm$  SD for eight rats in each group.

<sup>a</sup>: *P* < 0.05 compared to control, <sup>b</sup>: *P* < 0.05 compared to SP, <sup>c</sup>: *P* < 0.05 compared to CdCl<sub>2</sub>.

Abbreviations; TGF- $\beta$ , Transforming growth factor-beta; TNF- $\alpha$ , Tumor necrosis factor- $\alpha$ ; IL-10, Interleukin-10; IL-1 $\beta$ , Interleukin-1 $\beta$ ; SP, *Spirulina platensis*; CdCl<sub>2</sub>, Cadmium chloride.


The lobes of the gland were separated by connective tissue septa. (Fig. 1A and 1B).

Stained sections by H&E of the submandibular gland from the CdCl<sub>2</sub> group showed massive histological changes in the acini, as they degenerated, disarranged with disturbance of their architecture, showing widely cellular detached by connective tissue that contained dilated congested blood vessels (Fig. 1C and 1D). Some intralobular ducts displayed disarrangement of the lining epithelial cells with hyperchromatic nuclei and fatty infiltration (Fig. 1E and 1F). While the CdCl<sub>2</sub>/SP group showed normal acinar structure with the normal arrangement of acini and ducts with increasing in their number (Fig. 1G).

Histochemical examination of PAS-stained sections of the submandibular salivary gland in the control and SP groups showed that most of the acinar cells contain a mixture of both neutral and acid polysaccharides as they have deep purple color, while the duct system showed faint PAS-positive reaction (Fig. 2A and 2B).

Stained sections of the submandibular gland by PAS from the CdCl<sub>2</sub> group showed a marked decrease in the number of neutral mucopolysaccharides as there was a decrease in the purple staining in comparison to that observed in control sections (Fig. 2C). The CdCl<sub>2</sub>/SP group revealed normal PAS + ve acinar cells due to an increase in the content of mucopolysaccharides (Fig. 2D).





**Fig. 1.** A) A photomicrograph of rat submandibular gland from the control group showing many lobules (circle) containing mainly serous acini (s) and intralobular ducts (d). The intercalated ducts are lined by cuboidal epithelium. Notice connective tissue septa (c) between lobules and contain blood vessels (v) and interlobular space (l). H&E, microscopic magnification  $\times 200$ . B) Higher magnification photomicrograph of rat submandibular gland from the control group showing dark serous acinar cells (s) and pale mucous cells (m). Notice that some mucous acini are capped by serous demilunes (D), the myoepithelial cells are seen as related to the serous acini (arrow). H&E, microscopic magnification  $\times 400$ . C) A photomicrograph of rat submandibular gland from group II showing acinar architecture (serous and mucous). Notice that the intercalated ducts are lined by cuboidal epithelium (d) and congested blood vessels (v) between the acini. H&E, microscopic magnification  $\times 200$ . D) A photomicrograph of rat submandibular gland from group III showing acinar degeneration (arrows). Notice that the inflammatory cells infiltration (i), dilated congested blood vessels (V), and wide separation between the lobules (W). H&E, microscopic magnification  $\times 200$ . E) A photomicrograph of rat submandibular gland group III showing congested blood vessels (v) and ductal degeneration (d). Notice the acinar nuclear metachromatic (arrow). H&E, microscopic magnification  $\times 400$ . F) A photomicrograph of rat submandibular gland from The CdCl<sub>2</sub> group showing normal acinar cells (arrow). Notice that congested blood vessels (v) and normal duct (d) H&E, microscopic magnification  $\times 200$ . G) A photomicrograph of rat submandibular gland from The CdCl<sub>2</sub>/SP group showing both normal mucous and serous acini (arrow). H&E, microscopic magnification  $\times 400$ . H) Mean area percentage of histological score.

### 3.4. SP alleviated CdCl<sub>2</sub>-induced submandibular apoptotic, and morphometric changes in the rats

Immunostaining with BAX was mildly positive in control and SP glandular tissue (Fig. 3A and 3B); in the CdCl<sub>2</sub> group, it was strongly positive (Fig. 3C). In the CdCl<sub>2</sub>/SP group, the BAX immunostaining was mild to moderate (Fig. 3D).

The histological grading scale findings are represented in Fig. 1H in which there were significantly high ( $P < 0.001$ ) total points in CdCl<sub>2</sub> group obtained sections as compared to that of the control while significantly decreased ( $P < 0.001$ ) in the CdCl<sub>2</sub>/SP group when compared to CdCl<sub>2</sub> group.

The mean area percentages of PAS reaction for all groups were in Fig. 2E. There was a significant decrease ( $P < 0.05$ ) in PAS reaction in the CdCl<sub>2</sub> group compared with the control group. There was a significant increase in the mean area of staining cells ( $P < 0.001$ ) in the BAX reaction in the CdCl<sub>2</sub> group cells of the submandibular gland compared

with the control, whereas the BAX reaction in the CdCl<sub>2</sub>/SP group showed a significant decrease in positive staining cells ( $P < 0.001$ ) (Fig. 3E) when compared with the CdCl<sub>2</sub> group.


## 4. Discussion

The toxic effects of Cd have been described in many researches. Furthermore, the results of previous many *in vivo* animal studies have shown that repeated exposure to Cd can have detrimental effects on the salivary glands [29–31].

To the best of our knowledge, the effects of SP on Cd-induced submandibular biochemical and histopathological alterations have not been reported yet. The current results indicated that Cd-induced inflammatory and oxidative stress processes in the submandibular salivary glands, which also have been mirror-imaged in histopathology and immunohistochemistry. However, SP protected against these changes.

The Cd-treated group revealed decreased body and submandibular





**Fig. 2.** Photomicrographs of rat submandibular gland in A) The control group showing intense PAS stain in acinar cells and duct (arrow). B) The SP group shows intense PAS stain in acinar cells and ducts. C) The CdCl<sub>2</sub> group showed a marked decrease in the staining intensity of the secretory granules. Notice that increased numbers of intercalated ducts with decreased intensity of PAS stain (arrow). D) The CdCl<sub>2</sub>/SP group showed a moderate decrease in PAS staining acini (arrow). PAS, microscopic magnification  $\times 400$ . E) Mean area percentage of PAS staining.

glands' weight, according to previous findings [32,33]. This may be due to inhibition of the activity of digestive enzymes and decreased appetite induced by Cd [34,35].

In our study, Cd caused oxidative stress verified by increased MDA levels along with decreased catalase, SOD, GSH, and TAC.

In agreement with our results, several studies reported that Cd causes the inactivation of most of the antioxidant enzymes as it binds to the active sites of the sulfhydryl-group-containing enzymes resulting in the release of ROS, increased MDA, which is a product of lipid peroxidation, weakening of antioxidant defenses and DNA damage [36–38].

The potential interaction of Cd with the thiol groups present in the SOD protein has been observed. Moreover, Cd can exert inhibition on the other cofactors of SOD, ultimately leading to enzyme inactivation. As a vital antioxidant enzyme, CAT possesses heme as its active site prosthetic group. It has been noted that Cd can impede iron absorption and hinder the heme biosynthesis process [39].

In the current study, Cd caused elevation in the levels of TGF- $\beta$ , TNF- $\alpha$ , and IL-1 $\beta$ , however, the IL-10 level was reduced. These findings are matched with previous results which stated that Cd could increase the pro-inflammatory factor IL-1 $\beta$  and decrease the anti-inflammatory factor IL-10 [40].

IL-1 $\beta$  and TNF- $\alpha$ , play a vital role beginning of the inflammatory cascades that control the expression of other cytokines and chemokines [41].

It was reported that Cd induced the production of TNF- $\alpha$  from macrophages by inhibiting mitogen-activated protein kinase dephosphorylation [42]. The cytokine TNF- $\alpha$  serves as an enhancer and pro-longer of the inflammatory response through its activation of additional

cells to secrete both ROS and cytokines such as interleukin-1 $\beta$ , thereby promoting further inflammation and damage to the affected tissue [43].


It is proposed that exposure to Cd instigates the production of cytokines, thereby leading to disruptions in the cellular immune response [44].

The histopathological results in the current study agree with previous studies [25,29,45], which reported the toxic effects of Cd on the salivary glands with decreased intensity of the PAS reaction. The pathological changes in the morphology of the salivary gland tissue could be explained as Cd can cause an imbalance of the oxidative/antioxidative status along with triggered lipid peroxidation and oxidative protein changes [30].

In the present study, Cd caused an increased Bax immunostaining in the submandibular glands. Cd is supposed to cause disturbances in mRNA expression of the Bax gene which is involved in apoptosis [46]. The expression of Bax is augmented by Cd, leading to an overabundance of Ca<sup>2+</sup> in the mitochondria and an increase in the permeability of the transition pores. This, in turn, induces mitochondrial swelling, wherein the outer membranes of the mitochondria collapse and their contents exit into the cytoplasm, ultimately triggering the initiation of apoptosis [47].

Various experimental models have studied the role of antioxidants and dietary supplements in alleviating the toxic effects induced by Cd with documented positive outcomes in many of them [48–50].

The current study demonstrates that the administration of SP exhibited a reduction in the detrimental impact of Cd on the submandibular glands. Through co-administration of SP, a decrease in MDA levels was observed in addition to elevated levels of catalase, SOD, GSH,



**Fig. 3.** Photomicrographs of rat submandibular gland in A) The control group showing moderate BAX reaction in both acini and ducts (arrows). B) The SP group showed moderate BAX reaction in both acini and ducts (arrows). C) The CdCl<sub>2</sub> group showing a strong positive BAX reaction in acini and ducts (arrow). D) The CdCl<sub>2</sub>/SP group showing mild to moderate BAX reaction in acini and duct (arrow). Immunostaining for BAX microscopic magnification  $\times 400$ . E) Mean area percentage of BAX immunopositive cells.

and TAC in comparison to the Cd-treated group. Furthermore, SP resulted in a reduction of TGF- $\beta$ , TNF- $\alpha$ , and IL-1 $\beta$  levels, while increasing IL-10 levels. SP displayed amelioration of the histopathological and immunohistochemical alterations induced by Cd.

SP has been reported to possess significant benefits in combating various oxidative stress conditions initiated by toxic agents [51–53]. The potential protective effects exhibited by SP may be attributed to its abundant presence of active components with antioxidant properties such as proteins, phycocyanin, tocopherols, phenolic acids,  $\beta$ -carotenes, vitamins and minerals [54,55].

SP can induce powerful antioxidant enzymatic activities as well as prevention of lipid peroxidation, DNA damage, and free radicals

scavenging [56].

SP and its active compound, phycocyanin, show immunomodulatory effects as they can trigger the antibodies production and regulate the expression of genes responsible for cytokines and interleukin production [57].

## 5. Conclusion

Cadmium elicited damage in the submandibular salivary glands as verified by the biochemical, histopathological, and immunohistochemical changes. The present study has shown for the first time the protective effects of simultaneous administration of SP against these effects.



## CRediT authorship contribution statement

**Eman Mohamed Faruk:** Formal analysis, Conceptualization. **Fatma Ibrahim:** Writing – original draft, Investigation. **Dalia Ibrahim El-wafaei:** Visualization, Validation. **Yasmeen Mohammed Ismail El Sayed:** Resources, Data curation. **Mohamed Ghazy Attia Hablas:** Data curation, Validation. **Mahmoud M. Hassan:** Formal analysis. **Ahmed Mohammed Zaazaa:** Resources. **Kamal Mostafa Kamal:** Supervision.

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Data availability

The data that has been used is confidential.

## References

- [1] G. Genchi, M.S. Sinicropi, G. Lauria, et al., The effects of cadmium toxicity, *Int. J. Environ. Res. Public Health* 17 (2020) 3782.
- [2] H. Zhang, M. Reynolds, Cadmium exposure in living organisms: A short review, *Sci. Total Environ.* 678 (2019) 761–767.
- [3] H. Ali, E. Khan, I. Ilahi, Environmental chemistry and ecotoxicology of hazardous heavy metals: environmental persistence, toxicity, and bioaccumulation, *J. Chem.* 2019 (2019).
- [4] J.I. Anetor, Rising environmental cadmium levels in developing countries: threat to genome stability and health, *Niger. J. Physiol. Sci.* 27 (2012) 103–115.
- [5] M.R. Rahimzadeh, M.R. Rahimzadeh, S. Kazemi, et al., Cadmium toxicity and treatment: An update, *Caspian J. Intern. Med.* 8 (2017) 135.
- [6] W. Maret, J.-M. Moulis, The bioinorganic chemistry of cadmium in the context of its toxicity, *Cadmium: Toxicity Essentiality* (2013:) 1–29.
- [7] T. Moiseenko, N. Gashkina, Biogeochemistry of cadmium: anthropogenic dispersion, bioaccumulation, and ecotoxicity, *Geochem. Int.* 56 (2018) 798–811.
- [8] A. Menke, P. Muntner, E.K. Silbergeld, et al., Cadmium levels in urine and mortality among US adults, *Environ. Health Perspect.* 117 (2009) 190–196.
- [9] F. Thévenod, W.-K. Lee, Cadmium and cellular signaling cascades: interactions between cell death and survival pathways, *Arch. Toxicol.* 87 (2013) 1743–1786.
- [10] E.A. Belyaeva, D. Dymkowska, M.R. Więkowski, et al., Mitochondria as an important target in heavy metal toxicity in rat hepatoma AS-30D cells, *Toxicol. Appl. Pharmacol.* 231 (2008) 34–42.
- [11] A.A. Tinkov, T. Filippini, O.P. Ajsuvakova, et al., Cadmium and atherosclerosis: A review of toxicological mechanisms and a meta-analysis of epidemiologic studies, *Environ. Res.* 162 (2018) 240–260.
- [12] D. Tibullo, G.L. Volti, C. Giallongo, et al., Biochemical and clinical relevance of alpha lipoic acid: antioxidant and anti-inflammatory activity, molecular pathways and therapeutic potential, *Inflamm. Res.* 66 (2017) 947–959.
- [13] A. El-Baky, H. Hanaa, F. El Baz, et al., Enhancement of antioxidant production in *Spirulina platensis* under oxidative stress, *Acta Physiol. Plant.* 31 (2009) 623–631.
- [14] C.S. Ku, Y. Yang, Y. Park, et al., Health benefits of blue-green algae: prevention of cardiovascular disease and nonalcoholic fatty liver disease, *J. Med. Food* 16 (2013) 103–111.
- [15] M.N. Bin-Jumah, A.A. Al-Huqail, N. Abdelnaeim, et al., Potential protective effects of *Spirulina platensis* on liver, kidney, and brain acrylamide toxicity in rats, *Environ. Sci. Pollut. Res.* 28 (2021) 26653–26663.
- [16] D. Banji, O.J. Banji, N.G. Pratusha, et al., Investigation on the role of *Spirulina platensis* in ameliorating behavioural changes, thyroid dysfunction and oxidative stress in offspring of pregnant rats exposed to fluoride, *Food Chem.* 140 (2013) 321–331.
- [17] M. Afkhami-Ardakani, S. Hasanazadeh, R. Shahrooz, et al., Antioxidant effects of *Spirulina platensis* (Arthrospira platensis) on cyclophosphamide-induced testicular injury in rats, in: *Veterinary Research Forum*, Faculty of Veterinary Medicine, Urmia University, Urmia, Iran, 2018, p. 35.
- [18] K. Ahmed, R. Korany, H. El Halawany, et al., *Spirulina platensis* alleviates arsenic-induced toxicity in male rats: biochemical, histopathological and immunohistochemical studies, *Adv. Anim. Vet. Sci.* 7 (2019) 701–710.
- [19] M. Gargouri, A. Soussi, A. Akrouti, et al., Ameliorative effects of *Spirulina platensis* against lead-induced nephrotoxicity in newborn rats: modulation of oxidative stress and histopathological changes, *EXCLI J.* 17 (2018) 215.
- [20] G.E. El-Desoky, S.A. Bashandy, I.M. Alhazza, et al., Improvement of mercuric chloride-induced testis injuries and sperm quality deteriorations by *Spirulina platensis* in rats, *PLoS One* 8 (2013) e59177.
- [21] F. Ibrahim, M.A. Nomier, L.M. Sabik, et al., Manganese-induced neurotoxicity and the potential protective effects of lipoic acid and *Spirulina platensis*, *Toxicol. Mech. Methods* 30 (2020) 497–507.
- [22] M.A. Ibrahim, A.H. Almaeen, M. Abd El Moneim, et al., Cadmium-induced hematological, renal, and hepatic toxicity: the amelioration by *Spirulina platensis*, *Saudi J. Forensic Med. Sci.* 1 (2018) 5.
- [23] N. Simsek, A. Karadeniz, Y. Kalkan, et al., *Spirulina platensis* feeding inhibited the anemia-and leucopenia-induced lead and cadmium in rats, *J. Hazard. Mater.* 164 (2009) 1304–1309.
- [24] A.M. Attia, F.A. Ibrahim, N.A. Abd El-Latif, et al., Antioxidant effects of curcumin against cadmium chloride-induced oxidative stress in the blood of rats, *J. Pharmacogn. Phytother.* 6 (2014) 33–40.
- [25] L.E. Amin, W. Zedan, M.T. Abou Elkhier, et al., Cadmium induced apoptosis and cellular toxicity in rats major salivary glands, *Egypt. Dent. J.* 62 (2016) 653–660.
- [26] J.D. Bancroft, C. Layton, Connective and other mesenchymal tissues with their stains, *Bancroft's Theory Pract. Histol. Tech. E-Book* 153 (2018).
- [27] J. Kiernan, *Histological and histochemical methods*, Scion Publishing Ltd, 2015.
- [28] J. Cuzick, A Wilcoxon-type test for trend, *Stat. Med.* 4 (1985) 87–90.
- [29] E.H. Ahmed, M.E. Grawish, M.M. Anees, et al., Impact of bone marrow mesenchymal stem cells on the submandibular gland structure of adult male albino rats exposed to cadmium chloride toxicity, *Arch. Oral Biol.* 145 (2023) 105585.
- [30] B.M. Onopiuk, Z.N. Dąbrowska, J. Rogalska, et al., The beneficial impact of the black chokeberry extract against the oxidative stress in the sublingual salivary gland of rats intoxicated with cadmium, *Oxid. Med. Cell. Longev.* 2021 (2021).
- [31] M. Abdollahi, A. Bahreini-Moghadam, B. Emami, et al., Increasing intracellular cAMP and cGMP inhibits cadmium-induced oxidative stress in rat submandibular saliva, *Comp. Biochem. Physiol. C: Toxicol. Pharmacol.* 135 (2003) 331–336.
- [32] J. Fang, S. Xie, Z. Chen, et al., Protective effect of vitamin E on cadmium-induced renal oxidative damage and apoptosis in rats, *Biol. Trace Elem. Res.* (2021) 1–13.
- [33] S. Asagba, M. Adaike, H. Kadiri, et al., Influence of aqueous extract of *Hibiscus sabdariffa* L. petal on cadmium toxicity in rats, *Biol. Trace Elem. Res.* 115 (2007) 47–57.
- [34] G. Eriyamremu, S. Asagba, E. Onyeneke, et al., Changes in carboxypeptidase A, dipeptidase and Na<sup>+</sup>/K<sup>+</sup> ATPase activities in the intestine of rats orally exposed to different doses of cadmium, *Biometals* 18 (2005) 1–6.
- [35] J. Wang, H. Zhu, S. Lin, et al., Protective effect of naringenin against cadmium-induced testicular toxicity in male SD rats, *J. Inorg. Biochem.* 214 (2021) 111310.
- [36] M. Hormozi, R. Mirzaei, A. Nakhaee, et al., The biochemical effects of occupational exposure to lead and cadmium on markers of oxidative stress and antioxidant enzymes activity in the blood of glaziers in tile industry, *Toxicol. Ind. Health* 34 (2018) 459–467.
- [37] P. Kostecka-Sochoń, B.M. Onopiuk, E. Dąbrowska, Protective effect of increased zinc supply against oxidative damage of sublingual gland in chronic exposure to cadmium: experimental study on rats, *Oxid. Med. Cell. Longev.* 2018 (2018).
- [38] D. Quig, Cysteine metabolism and metal toxicity, *Altern. Med. Rev.* 3 (1998) 262–270.
- [39] A. Schauder, A. Avital, Z. Malik, Regulation and gene expression of heme synthesis under heavy metal exposure-review, *J. Environ. Pathol. Toxicol. Oncol.* 29 (2010).
- [40] X. Zhao, S. Wang, X. Li, et al., Cadmium exposure induces TNF- $\alpha$ -mediated necroptosis via FPR2/TGFR- $\beta$ /NF- $\kappa$ B pathway in swine myocardium, *Toxicology* 453 (2021) 152733.
- [41] M. Låg, D. Rodionov, J. Øvrevik, et al., Cadmium-induced inflammatory responses in cells relevant for lung toxicity: Expression and release of cytokines in fibroblasts, epithelial cells and macrophages, *Toxicol. Lett.* 193 (2010) 252–260.
- [42] H. Haase, J.L. Ober-Blobaum, G. Engelhardt, et al., Cadmium ions induce monocytic production of tumor necrosis factor- $\alpha$  by inhibiting mitogen activated protein kinase dephosphorylation, *Toxicol. Lett.* 198 (2010) 152–158.
- [43] M.I. Elmallah, M.F. Elkhadragey, E.M. Al-Olayan, et al., Protective effect of *Fragaria ananassa* crude extract on cadmium-induced lipid peroxidation, antioxidant enzymes suppression, and apoptosis in rat testes, *Int. J. Mol. Sci.* 18 (2017) 957.
- [44] A.E. Turley, J.W. Zagorski, R.C. Kennedy, et al., Chronic low-level cadmium exposure in rats affects cytokine production by activated T cells, *Toxicol. Res.* 8 (2019) 227–237.
- [45] E.S. Elroghy, R.I. Yassien, Role of montelukast in alleviation of toxicity induced by cadmium in thyroid gland of adult male albino rats. (Histological, Immunohistochemical and Biochemical Study), *Egyptian J. Histol.* 44 (2021) 979–992.
- [46] A. Abdeen, O.A. Abou-Zaid, H.A. Abdel-Maksoud, et al., Cadmium overload modulates piroxicam-regulated oxidative damage and apoptotic pathways, *Environ. Sci. Pollut. Res.* 26 (2019) 25167–25177.
- [47] Y. Yuan, C.-y. Jiang, H. Xu, et al., Cadmium-induced apoptosis in primary rat cerebral cortical neurons culture is mediated by a calcium signaling pathway, *PLoS One* 8 (2013) e64330.
- [48] V. Unsal, T. Dalkiran, M. Çiçek, et al., The role of natural antioxidants against reactive oxygen species produced by cadmium toxicity: a review, *Adv. Pharm. Bull.* 10 (2020) 184.
- [49] B.E. Oyinloye, B.O. Ajiboye, O.A. Ojo, et al., Cardioprotective and antioxidant influence of aqueous extracts from Sesamum indicum seeds on oxidative stress induced by cadmium in wistar rats, *Pharmacogn. Mag.* 12 (2016) S170.
- [50] M.A. Ibrahim, A.H. Almaeen, M. Abd El Moneim, et al., Cadmium-induced hematological, renal, and hepatic toxicity: the amelioration by: *Spirulina platensis*, *Saudi J. Forensic Med. Sci.* 1 (2018) 5–13.
- [51] R. Korany, K.S. Ahmed, H. Halawany, et al., Effect of long-term arsenic exposure on female Albino rats with special reference to the protective role of *Spirulina platensis*, *Explor. Anim. Med. Res.* 9 (2019) 125–136.
- [52] N.Z. Eleiwa, A.A. Galal, R.M. Abd El-Aziz, et al., Antioxidant activity of *Spirulina platensis* alleviates doxorubicin-induced oxidative stress and reprotoxicity in male rats, *Orient. Pharm. Exp. Med.* 18 (2018) 87–95.
- [53] M.R. Hassanen, M.K. Mahfouz, A.S. Farid, et al., Biochemical effects of *Spirulina platensis* against oxidative stress caused by doxorubicin, *Benha Veterinary Med. J.* 28 (2015) 147–154.

- [54] N. Avdagić, E. Ćosović, E. Nakaš-Ićindić, et al., Spirulina platensis protects against renal injury in rats with gentamicin-induced acute tubular necrosis, *Bosn. J. Basic Med. Sci.* 8 (2008) 331.
- [55] Z. Khan, P. Bhadouria, P. Bisen, Nutritional and therapeutic potential of Spirulina, *Curr. Pharm. Biotechnol.* 6 (2005) 373–379.
- [56] N.K. Abdelkhalek, E.W. Ghazy, M.M. Abdel-Daim, Pharmacodynamic interaction of Spirulina platensis and deltamethrin in freshwater fish Nile tilapia, *Oreochromis niloticus*: impact on lipid peroxidation and oxidative stress, *Environ. Sci. Pollut. Res.* 22 (2015) 3023–3031.
- [57] Q. Wu, L. Liu, A. Miron, et al., The antioxidant, immunomodulatory, and anti-inflammatory activities of Spirulina: an overview, *Arch. Toxicol.* 90 (2016) 1817–1840.