

The relationship between distance of breast cancer from the skin and incidence of axillary nodal metastasis in female patients with early cancer breast: correlation between radiological and pathological distance

Mohamed S Essa,¹ Mohamed H Faheem,² Rana Abdalla,³ Mohamed E Zayed¹

¹Department of General Surgery, Benha University Hospital, Faculty of Medicine, Benha University, Benha, Egypt

²Department of Radiology, Benha University Hospital, Faculty of Medicine, Benha University, Benha, Egypt

³Department of pathology, Faculty of Medicine, Benha University, Benha, Egypt

Corresponding author: Mohamed
Elsayed Zayed

Department of General Surgery,
Faculty of Medicine, Benha University,
Benha, Egypt

E mail:
drmohamedelsaid1983@gmail.com

Mobile: 01110243330

Background: The aim of this study is to evaluate the relation between proximity of breast cancer to the skin and incidence of axillary nodal metastasis in order to clarify a new guideline in treatment of early cancer breast.

Methods: This study includes 50 female patients with early cancer breast (T1 and T2). All patients underwent breast ultrasonography (U/S) with special confirmation on the breast cancer distance from skin surface (radiological distance) in addition to pathological assessment of the distance (pathological distance) after surgical excision and its correlation with radiological distance. Breast conservative surgery (BCS) with axillary clearance was done for 46 patients, while 4 patients underwent modified radical mastectomy (MRM).

Results: This study showed that the more proximal the cancer from the skin, the higher the incidence of axillary lymph node (LN) metastasis, and the cut off radiological distance was 1.55cm while cut off pathological distance was 1.5 cm. It also proved that U/S is an accurate and reliable method in assessing the breast cancer distance.

Conclusion: Our results revealed that breast cancers located closer to the skin surface have a higher incidence of axillary LN metastasis.

Keywords: Early cancer breast cancer, Distance from the skin surface, Axillary lymph node metastasis, Ultrasonography in cancer breast.

Background

In spite of pathological features and molecular biology of cancer breast, the status of axillary lymph nodes is one of the most reliable and important prognostic factors in addition to its role in guidance of adjuvant cancer breast treatment.¹

Axillary LN dissection (ALND) has been a routine part of cancer breast treatment for long time because it provides data about the status of axillary LN (prognostic role) in addition to removal of axillary tumors in patients presented with positive LNs (therapeutic role). Incidence of postoperative complications of ALND are 15–30% and include postoperative bleeding, local swelling, numbness, decrease range of motion, wound infection, neuropathy, and chronic lymphedema.² So predication of axillary nodal status before surgery is essential to prepare patient psychologically for ANLD and in planning for immediate breast reconstruction in addition to avoidance of ANLD if axilla is negative for metastasis.³

Several studies have reported that the lymphatic pathway present in the dermis is responsible for cancer breast metastatic pathway and suggest that the parenchyma of the breast is not rich in lymphatic plexus than that in the superficial dermal and subdermal layers.⁴

Only few studies have investigated the relation between the breast cancer distance from skin surface and axillary nodal deposits. U/S is a more accurate tool for detection of the distances from the skin. Distances of cancer breast from the skin on mammography are subject to variability because of

patient positioning.^{5,6} The aim of our study is to evaluate the relation between breast cancer distance from the skin and axillary LN metastasis

Methods

Study Design:

This prospective study includes 50 female patients with early cancer breast who were diagnosed and managed at General Surgery Department, Benha University Hospital from April 2017 to October 2019.

Inclusion criteria:

- 1-Female patients with T1 or T2 cancer breast.
- 2-Patients who have tumors visible on breast U/S or mammography.

Exclusion criteria:

- 1-Locally advanced breast cancer (T3, T4).
- 2-Metastatic cancer breast.
- 3-Multicentric breast cancer.
- 4-Benign breast lumps.
- 5-Women who received neoadjuvant chemotherapy.

After approval of the study by Benha Faculty of Medicine Ethical Committee and obtaining written informed consent

from the patients for the participation in the study. Patients were fully informed about the hazards and benefits of the surgery. The Patients assessed by a multidisciplinary team (includes one or more specialized representative from general surgery, pathology, radiology, radiotherapy, and medical oncology) and patients enrolled in the study if they fulfilled our inclusion criteria. All patients underwent the following:

1-Full detailed history.

2-Clinical examination.

3-Laboratory investigations: Complete blood picture (CBC), fasting and postprandial blood glucose liver function tests, renal function tests.

4-Radiological investigations (measurement of cancer distance from the skin).

Both breasts were examined by radiologist using US device equipped with linear transducer for three-dimensional image (5-12MHz). In all patients the distance was measured perpendicularly from the skin surface to the anterior hypoechoic edge of the tumor before the biopsy. All US images are recorded and subsequently taken for retrospective measurements of breast cancer proximity from surface of the skin (**Fig. 1**). Bilateral mammography also done.




Fig 1: Distance of cancer breast from the skin by U/S (white line).

5-Metastatic work up: Computed tomography (CT) of chest and abdomen in addition to bone scan if indicated.

6-Pathological evaluation (measurement of cancer distance from the skin)

Wide local excision (WLE) of the breast lump, is done with an elliptical incision of the overlying skin then it is sent for histopathological assessment with measurement of pathological distance (**Fig. 2**). The specimen is oriented by stitches to detect the margins for further management.

Operative plan:

Breast conservative surgery (BCS) or modified radical mastectomy (MRM) was explained to the patient preoperatively and informed consent was taken. Type of surgery done for our patients were conducted based on type of histopathology, size of the mass, breast size, site of the mass and socioeconomic status. Patients in our study underwent either BCS with axillary clearance or MRM (**Fig. 2**).




Fig 2: BCS (A) elliptical incision, (B) Orientation of specimen, (C) Mobilization of breast pillar, (D) Suturing of pillars, (E) Axillary clearance and (F) Suturing of the skin.

A final pathology report is included with special emphasis on the safety margins, histopathological type, distance from the skin and number of involved axillary LNs in addition to the immunohistochemical assay for oestrogen, progesterone receptors status (ER and PR) and Her2/neu expression in the specimen. In case of BCS, if the safety margin is inadequate (<1 cm) redo excision is done.

After fulfillment of the patients' data, they are informed regarding the dates of the regular follow-up and they were referred to the outpatient oncology clinic for further management plan regarding radiotherapy, chemotherapy in addition to hormonal therapy according to the final pathology report.

Primary endpoint:

Evaluation of the prognostic importance of assessing the proximity of cancer breast to the skin and incidence of axillary LN metastasis.

Secondary outcomes:

Evaluating the sensitivity of breast U/S and pathological assessment in detecting the distance between the cancer breast and the skin.

Statistical analysis:

Data were statistically described in terms of mean \pm standard deviation (\pm SD), median and range, or frequencies (number of cases) and percentages when appropriate. Comparison of numerical variables between the study groups was done using Kruskal Wallis test. Within group comparison of distance was done using paired t test. For comparing categorical data, Chi-square (χ^2) test was performed. Exact test was used instead when the expected frequency is less than 5. Accuracy was represented using the terms sensitivity, and specificity. Receiver operator characteristic (ROC) analysis was used to determine the optimum cut off value for the studied diagnostic markers. p values less than 0.05 was considered statistically significant. All statistical calculations were done using computer program IBM SPSS (Statistical Package for the Social Science; IBM Corp, Armonk, NY, USA) release 22 for Microsoft Windows.

Results

This prospective study includes 50 female patients presented with early cancer breast (T1 and T2). BCS was done for 46 patients, while 4 patients underwent MRM due to centrally located tumors, tumors that are large in relation to small breast size or sometimes patient wishes. There was no statistically significant difference between the age

of the patient (Mean \pm SD, 46.78 \pm 8.57) and incidence of axillary LN metastasis, (P= 0.165) (**Table 10**). There was no significant statistical variation between the location of tumor and the incidence of axillary nodal metastasis, (P= 0.175) (**Table 2**). The relation between US BIRAD classification of breast mass and incidence of axillary LN metastasis showed no statistically significant difference, (P= 0.486) (**Table 3**). There was no significant variation between tumor size (T stage) and incidence of nodal metastasis. (P= 0.758) (**Table 4**). Mean size of the tumor was 2.65cm. There was no statistically significant variation noted for ER, PR, or HER2 statuses and incidence of axillary nodal deposits, (P=0.267, P=0.124, P=0.542, respectively), (**Table 5,6**).

Primary outcome:

There was statistically significant variation between the incidence of axillary nodal metastasis and the radiological distance, (p value < 0.001). ROC analysis revealed that the most accurate cut off radiological distance was at 1.55cm that achieved sensitivity and specificity of 100% and 72.7, respectively. This means that early cancer breast with distance \leq 1.5 cm from the skin are more likely to develop axillary LN metastasis, and those with tumors that are deep to this cut off level are less likely to have axillary nodal metastasis. There was statistically significant difference between the incidence of axillary nodal deposits and the pathological distance, (P < 0.001), ROC analysis revealed that the most accurate cut off pathological distance was at 1.5 cm that achieved 100% sensitivity and 78.6% specificity (**Fig. 3**).

17 patients of 50 had positive axillary LNs, 8 of them were

staged as N1 and the other 9 patients were staged as N2. There was a significant statistical difference between the radiological distance and progression of axillary nodal metastasis from N1 to N2 stage, (P< 0.014), ROC analysis showed that 1.35 cm radiological distance was the most accurate cut off distance that achieved 100% sensitivity and 60% specificity. There was a significant statistical variation between pathological distance and N1 to N2 progression of axillary LN metastasis, (P< 0.001), ROC analysis revealed that 1.25 cm pathological distance achieved 83.3% sensitivity and 100% specificity, and 1.4 cm achieved 86.3 % sensitivity and 66.7% Specificity.

By using the T-test the mean radiological distance was 1.46 cm, while the mean pathological distance was 1.59cm. The mean difference between both measurements was only 0.13cm. There was no significant statistical difference between the radiological and pathological distance, (P = 0.647).

Secondary outcome:

U/S is an accurate and reliable method in a assessing the breast cancer distance, ROC analysis also showed that the overall accuracy of U/S in this test is 88.6%.




Fig 3: ROC analysis (A) radiological distance, (B) pathological distance.




Fig 4: ROC analysis of axillary nodal metastasis progression from N1 to N2 (a) radiological distance, (b) pathological distance.

Table 1: Relation between patient's age and axillary nodal metastasis

N		Age
0	Number	33
	Mean	51.87
	SD	6.483
1	Median	51.00
	Number	8
	Mean	46.78
2	SD	4.734
	Median	46.50
	Number	9
Total	Mean	52.67
	SD	8.570
	Median	52.00
Total	Number	50

SD: Standard deviation.

Table 2: Relation between breast tumor location and axillary nodal metastasis

		0	N		Total	
			1	2		
Tumor location	Central	Count	4	0	0	4
		% within tumor quadrant	100.0%	0.0%	0.0%	100.0%
		% within N	12.1%	0.0%	0.0%	8.0%
	Lower inner quadrant	Count	2	4	1	7
		% within tumor quadrant	28.5%	57.1%	14.2%	100.0%
		% within N	6.06%	50.0%	11.1%	14.0%
	Lower outer quadrant	Count	4	2	2	8
		% within tumor quadrant	50.0%	25.0%	25.0%	100.0%
		% within N	12.1%	25.0%	22.2%	16.0%
	Upper inner quadrant	Count	3	2	1	6
		% within tumor quadrant	50.0%	33.3%	16.6%	100.0%
		% within N	9.09%	25.0%	11.1%	12.0%
	Upper outer quadrant	Count	20	0	5	25
		% within tumor quadrant	80.0%	0.0%	20.0%	100.0%
		% within N	60.6%	0.0%	55.5%	50.0%
	Total	Count	33	8	9	50
		% within tumor quadrant	66.0%	16.0%	18.0%	100.0%
		% within N	100.0%	100.0%	100.0%	100.0%

Table 3: Relation between tumor BIRAD and axillary nodal metastasis

		0	N		Total	
			1	2		
BIRAD	4A	Count	2	0	0	2
		% within BIRAD	100.0%	0.0%	0.0%	100.0%
		% within N	6.3%	0.0%	0.0%	4.0%
	4B	Count	6	2	0	8
		% within BIRAD	75.0%	25.0%	0.0%	100.0%
		% within N	18.8%	22.2%	0.0%	16.0%
	4c	Count	11	1	4	16
		% within BIRAD	68.8%	6.3%	25.0%	100.0%
		% within N	34.4%	11.1%	44.4%	32.0%
	5	Count	13	6	5	24
		% within BIRAD	54.2%	25.0%	20.8%	100.0%
		% within N	40.6%	66.7%	55.6%	48.0%
	Total	Count	32	9	9	50
		% within BIRAD	64.0%	18.0%	18%	100.0%
		% within N	100.0%	100.0%	100.0%	100.0%

Table 4: Relation between breast tumor size (T) and axillary nodal metastasis

		0	N		Total	
			1	2		
T stage	1	Count	8	3	4	15
		% within T	53.3%	20.0%	26.7%	100.0%
		% within N	26.7%	30.0%	40.0%	30.0%
	2	Count	22	7	6	35
		% within T	62.9%	20.0%	17.1%	100.0%
		% within N	73.3%	70.0%	60.0%	70.0%
Total	Count	30	10	10	50	
	% within T	60%	20.0%	20.0%	100.0%	
	% within N	100.0%	100.0%	100.0%	100.0%	

Table 5: Relation between breast tumor ER and PR status and axillary nodal metastasis

		N			Total	
		0	1	2		
Estrogen receptor(ER)	Negative	Count	3	0	2	5
		% within ER	60.0%	0.0%	40.0%	100.0%
		% within N	9.1%	0.0%	22.2%	10.0%
	Positive	Count	30	8	7	45
		% within ER	66.7%	17.8%	15.6%	100.0%
		% within N	90.9%	100.0%	77.8%	90.0%
Total	Count	33	8	9	50	
	% within ER	66.0%	16.0%	18%	100.0%	
	% within N	100.0%	100.0%	100.0%	100.0%	
Progesterone receptor (PR)	Negative	Count	5	1	5	11
		% within PR	45.5%	9.1%	45.5%	100.0%
		% within N	15.6%	11.1%	55.6%	22.0%
	Positive	Count	27	8	4	39
		% within PR	69.2%	20.5%	10.3%	100.0%
		% within N	84.4%	88.9%	44.4%	87.0%
Total	Count	32	9	9	50	
	% within PR	64.0%	18.0%	18.0%	100.0%	
	% within N	100.0%	100.0%	100.0%	100.0%	

Table 6: Relation between breast tumor Her2/neu status and axillary nodal metastasis

		N			Total	
		0	1	2		
HER2	Negative	Count	33	8	5	46
		% within HER2	71.7%	17.4%	10.9%	100.0%
		% within N	91.7%	88.9%	100.0%	92.0%
	Positive	Count	3	1	0	4
		% within HER2	75.0%	25.0%	0.0%	100.0%
		% within N	8.3%	11.1%	0.0%	8.0%
	Total	Count	36	9	5	50
		% within HER2	72%	18.0%	10.0%	100.0%
		% within N	100.0%	100.0%	100.0%	100.0%

Discussion

Status of axillary LN is one of the most important and reliable prognostic indicators and is an essential factor to detect staging and treatment of cancer breast cancer.¹ Prediction of axillary nodal status before surgery is important for many reasons. It can prepare patients psychologically for ANLD if required in addition to it can assist with contingency planning for patients who desire immediate breast reconstruction. Accurate prediction of axillary LN status can avoid ANLD and its associated complications.^{7,8,9}

There are many clinical, histopathological, and molecular features have been related to the likelihood of axillary nodal metastases in patients with cancer breast.¹⁰⁻¹⁴ Several nomograms have been initiated for the prediction of axillary LN metastasis, such as those developed at Memorial Sloan Kettering Cancer Center (MSKCC) and MD Anderson Cancer Center. These nomograms have specified several predictors of axillary nodal deposits such as: size of the tumor, age of the patient, anatomical location, lymphovascular invasion (LVI), histological type of the tumor, tumor grading, hormonal status of the tumor (ER and PR), in addition to focality and centrality of the tumor. These developed nomograms did not involve breast cancer distance from the skin and nipple – areola complex in their validation.^{5,6,15}

In our study, 50 patients of early breast cancer (T1, T2) were involved in which they underwent BCS (46 patients) or MRM (4 patients), for all patients the radiological and pathological distances were measured. 33 patients (66%) were node negative and 17 patient (34%) were node positive, 8 (16%) and 9 (18%) patients staged as N1 and N2, respectively. ANLD was done for all patients and the mean number of excised LNs was 21.78. This compares well with the previous studies like, Cunningham et al., study that includes 209 patients, 61 (29%) had axillary nodal metastasis, and 148 (71%) had no known metastases and the study of Ansari et al. in which data were collected on 233 early breast cancer patients, of which 177 (76%) were node negative and 56 (24%) were node positive.^{4,5}

The mean age of the patients in our study was 51.87±6.48 for node negative patients, 46.78± 4.73 for N1 patient group, and 52.67±8.57 for N2 patient group. Eom et al., reported that the mean age for node negative patients is 51.07 group, and for node positive patients is 48.69 while Ansari et al., reported that the mean age of all patients was 66.2 years while the mean age for node negative and positive patients was 66.9 and 63.9, respectively.^{16,5} In our study there was no statistically significant difference between the age of the patient and the incidence of axillary nodal metastasis (P= 0.165). Cunningham et al., reported similar result and declared there was no statistical significant variation between the age of the patient and the incidence of axillary LN metastasis, (P= 0.399).⁴

In the current study there was no significant statistical variation between the location of the tumor and incidence of axillary LN deposits, (P=0.185). This compares well with the study of Cunningham et al., that revealed there was no significant statistical variation between the tumor location and incidence of axillary nodal metastasis, (P=0.138).⁴ However some studies like Ansari et al., showed that there was a significant statistical variation between the tumor location and incidence of axillary LN metastasis, (P=0.001). This may be explained by sample size difference.⁵

In our study there was no statistically significant variation between the tumor size and the incidence of axillary nodal metastasis, (P= 0.758). Mean tumour size in our study

was 2.65 cm. Eom et al., reported that the mean tumor size for node negative and positive patients was 1.92 and 2.76 cm, respectively and this matching with our result that showed there was no statistically significant relation between the tumor size (T) and incidence of axillary LN metastasis, (P=0.709).¹⁶ Other studies reported different data like Ansari et al., in which the mean tumor size for all patients was 1.34 cm and it was 1.24 cm and 1.68 for node negative and positive patients, respectively and showed significant statistical difference between the size of tumor and incidence of axillary nodal metastasis. Patients T1c and T2 stages were 7.7 and 8.6 times more likely to have positive nodes, respectively than patients with T1a and T1b stages, (P=0.001).⁵ Cunningham et al., reported that tumors in the node positive patients tended to be larger (P= 0.002).⁴

In the present study no significant statistical difference was observed for ER, PR, or HER2 status and axillary LN positivity, (P=0.267, 0.124, 0.542) respectively. This similar to the study reported by Eom et al. in which there is no significant variations were observed for ER, PR, or HER2 statuses between patients with positive and negative axillary LN (P=0.478, P=0.534, P=0.422) respectively.¹⁶ Ansari et al. showed also that there was no significant variation noted for ER, PR, or HER2 statuses between patients with positive and negative axillary nodes, (P=1.0, P=0.14, P=1.0) respectively.⁵

In the current study, it was found that there is a statistically significant correlation between the incidence of axillary LN metastasis and the radiological distance (P< 0.001), the most accurate cut off level at 1.55 cm achieved 100% sensitivity and 72.7% specificity. Also, it was noted that there was a statistically significant correlation between the incidence of axillary LN metastasis and the pathological distance (P< 0.001), the most accurate cut off at 1.5 cm achieved 100% sensitivity and 78.6% specificity. It was obvious in our study that no patient with cancer deep to 1.55 cm from the skin had any positive axillary LNs. This means that 1.55 cm is the most likely cut off distance at which positive axillary LNs is to be predicted. Eom et al., reported that patients with cancer breast closer to the skin are more likely to develop positive axillary LNs than those with tumor far from the skin, (P= 0.047).¹⁶ This also compared well with Ansari et al., who reported that closer proximity of the cancer to the skin was associated with positive axillary LNs and each 1 mm decrease in the cancer distance from the skin was associated with a 15% increased likelihood of lymph node metastasis.⁵ Ansari et al., studied also the relationship between the size of the largest metastatic axillary LN and the breast cancer distance from the skin and reported that there was no statistically significant association between the size of the largest LN metastasis and distance from the skin, (P= 0.72) and also there was no significant statistical correlation between the number of positive axillary LNs and the distance of the tumor from the skin, (P=0.29).⁵ Cunningham et al., reported that of none of the 26 cancers with distance >14 mm measured by U/S had metastasized to axillary LNs. In logistic regression analysis size of the tumor, histologic grade and proximity of the tumor to the skin were significantly associated with odds of axillary nodal metastasis. Among cancers within 14 mm distance, proximity was not an independent predictor.⁴

Our study revealed that there was a statistically significant relation between the radiological distance and progression of axillary LN metastasis from N1 to N2 stage, (P< 0.014) and the most accurate cut off radiologica distance at which progression of from N1 to N2 stage was most likely to occur at 1.35 cm that achieved 100% sensitivity and 60% Specificity. There was a significant statistical correlation between the

pathological distance and axillary LN metastasis progression from N1 to N2 stage, ($P < 0.001$) and the most accurate cut off pathological distance was between 1.25 cm that achieved 83.3% sensitivity and 100% specificity, and 1.4cm that achieved 86.3 % sensitivity and 66.7% specificity.

We also compared radiological distance with the pathological distance in order to clarify the accuracy of U/S in detecting breast cancer distance. We found that the mean radiological distance was 1.46cm, while the mean pathological distance was 1.59cm. The mean difference between both distances was only 0.17 mm. There was no significant statistical variation between the 2 methods of measurement, ($P=0.647$). The overall accuracy of the U/S was 88.6%. This means that U/S is considered to be a reliable investigating method detection of the distance between the breast cancer and the surface of the skin.

There is some limitations of this study that have to be addressed including technical factors that could have affected the distance from the skin i.e the scan technique and ability of the ultrasonography physician to obtain the proper image through the image plane that most accurately evaluate proximity of the tumor to the skin surface which may result in induced ultrasound measurement error. Some tumors had poorly defined edges, and others had hyperechoic halos surrounding the tumor, ultrasound characteristics that make accurate measurements are more difficult to obtain. The size of breast was not involved in this study, and the relation between tumor size and breast size may influence proximity to the skin, especially in smaller breasts and it is well known that distance from the skin is shorter in smaller breasts than in larger breasts; therefore, effect of breast size on nodal metastases cannot be excluded. Sample size of the patients was small which was also a limiting factor, so further prospective studies with larger patient numbers and a standardized US protocol would be required.

Ultimately, in the future, we may be looking to detect which patients are able to avoid any axillary surgery including sentinel lymph node biopsy (SLNB).

Conclusion

Cancer distance from the skin should be taking into consideration when preoperatively evaluating a patient with breast cancer before surgery and assessing the risk of axillary nodal deposits.

List of abbreviations:

U/S: Ultrasonography

ER: Estrogen receptor

PR: Progesterone receptor

ANLD: Axillary lymph node dissection

MKSCC: Memorial Sloan Kettering Cancer Center

LN: Lymph node

BIRAD: Breast Imaging Reporting and Data System

Declarations

Acknowledgements:

Not applicable

Compliance with Ethical Standards:

Conflict of interest

Mohamed S Essa: Declares that he has no conflict of interest.

Mohamed E Zayed: Declares that he has no conflict of interest.

Rana Abdalla: Declares that she has no conflict of interest.

Mohamed H Faheem: Declares that she has no conflict of interest.

Ethical approval

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

Consent to participate

Informed consent was obtained from all individual participants included in the study.

Consent for publication:

Written consent was obtained for publication of this study

Availability of data and materials:

The datasets used and/or analysed during the current study available from the corresponding author on reasonable request.

Funding:

The authors did not receive support from any organization for the submitted work.

Authors' contributions:

Mohamed S Essa: Study conception, design and Drafting of manuscript.

Mohamed E Zayed: Acquisition of data, analysis and interpretation of data.

Rana Abdalla: Acquisition of data and critical revision of manuscript.

Mohamed H Faheem: Acquisition of data and critical revision of manuscript

References

1. Colleoni M, Zahrieh D, Gelber RD, Holmberg SB, Mattsson JE, Rudenstam CM, et al. Site of primary tumor has a prognostic role in operable breast cancer: The international breast cancer study group experience. *J Clin Oncol.* 2005;23(7):1390–1400.
2. Rietman JS, Dijkstra PU, Geertzen JH, Baas P, De Vries J, Dolsma W, et al. Short term morbidity of the upper limb after sentinel lymph node biopsy or axillary lymph node dissection for Stage I or II breast carcinoma. *Cancer: Interdisciplinary International Journal of the American Cancer Society.* 2003;98(4):690-696.
3. Giuliano AE, Hunt KK, Ballman KV, Beitsch PD, Whitworth PW, Blumencranz PW, et al. Axillary dissection vs no axillary dissection in women with invasive breast cancer and sentinel node metastasis: a randomized clinical

- trial. *Jama*. 2011;305(6):569-575.
4. Cunningham JE, Jurj AL, Oman L, Stonerock AE, Nitscheva DK, Cupples TE. Is risk of axillary lymph node metastasis associated with proximity of breast cancer to the skin? *Breast cancer research and treatment*. 2006;100(3):319-328.
 5. Ansari B, Morton MJ, Adamczyk DL, Jones KN, Brodt JK, Degnim AC, et al. Distance of breast cancer from the skin and nipple impacts axillary nodal metastases. *Annals of surgical oncology*. 2011;18(11):3174-3180.
 6. Veerapong J, Boughey JC, Mittendorf EA, Harrell R, Bassett R, Ross M, et al. A validated risk assessment of sentinel lymph node involvement in breast cancer patients. *Ann Surg Oncol*. 2011;18:S59 (Abstract).
 7. Silverstein MJ, Skinner KA, Lomis TJ. Predicting axillary nodal positivity in 2282 patients with breast carcinoma. *World journal of surgery*. 2011;25(6):767-772.
 8. Giuliano AE, McCall L, Beitsch P, Whitworth PW, Blumencranz P, Leitch AM, et al. Locoregional recurrence after sentinel lymph node dissection with or without axillary dissection in patients with sentinel lymph node metastases: the American College of Surgeons Oncology Group Z0011 randomized trial. *Annals of surgery*. 2010;252(3):426.
 9. Suami H, Pan WR, Taylor GI. Historical review of breast lymphatic studies. *Clinical Anatomy: The Official Journal of the American Association of Clinical Anatomists and the British Association of Clinical Anatomists*. 2009;22(5):531-536.
 10. Rahusen FD, Torrenga H, Van Diest PJ, Pijpers R, Van Der Wall E, Licht J, et al. Predictive factors for metastatic involvement of nonsentinel nodes in patients with breast cancer. *Arch Surg*. 2001;136:1059-1063.
 11. Cavalli LR. Molecular markers of breast axillary lymph node metastasis. *Expert Rev Mol Diagn*. 2009;9:441-454.
 12. Lee JH, Kim SH, Suh YJ, Shim BY, Kim HK. Predictors of axillary lymph node metastases (ALNM) in a Korean population with T1-2 breast carcinoma: triple negative breast cancer has a high incidence of ALNM irrespective of the tumor size. *Cancer Res Treat*. 2010;42:30-36.
 13. Rivadeneira DE, Simmons RM, Christos PJ, Hanna K, Daly JM, Osborne MP. Predictive factors associated with axillary lymph node metastases in T1a and T1b breast carcinomas: analysis in more than 900 patients. *J Am Coll Surg*. 2000;191:1-6.
 14. Patani NR, Dwek MV, Douek M. Predictors of axillary lymph node metastasis in breast cancer: a systematic review. *Eur J Surg Oncol*. 2007;33:409-419.
 15. Bevilacqua JL, Kattan MW, Fey JV, Cody III HS, Borgen PI, Van Zee KJ. Doctor, what are my chances of having a positive sentinel node? A validated nomogram for risk estimation. *Journal of clinical oncology*. 2007;25(24):3670-3679.
 16. Eom YH, Kim EJ, Chae BJ, Song BJ, Jung SS. The distance between breast cancer and the skin is associated with axillary nodal metastasis. *Journal of surgical oncology*. 2015;111(7):824-828.