

**Draft Manuscript for Review**

"Nephroprotective Potential of Ginger, Diosmin, and Their Combination  
on Induced Diabetic Nephropathy in a Rat Model: Light and Electron  
Microscopic Observations"

Journal:	Egyptian Journal of Histology
Manuscript ID:	EJH-2407-2109 (R3)
Manuscript Title:	Nephroprotective Potential of Ginger, Diosmin, and Their Combination on Induced Diabetic Nephropathy in a Rat Model: Light and Electron Microscopic Observations

## Nephroprotective Potential of Ginger, Diosmin, and Their Combination on Induced Diabetic Nephropathy in a Rat Model: Light and Electron Microscopic Study

### Abstract

**Background:** Hyperglycemia is a chronic illness that is associated with diabetes. Patients with diabetes have been found to have many alterations in their renal functioning. The aim of this study was to evaluate the potential benefits of diosmin and ginger, either separately or in combination, for diabetic nephropathy that was experimentally produced.

**Materials and Methods:** Group I, ten rats stood as a control group. All rats (except control group) were given a single intraperitoneal dose of alloxan (150 mg/kg body weight) to induce type 2 diabetes. Group II consisted of ten diabetic rats. Other ten diabetic rats received 100 mg/kg diosmin orally for six weeks as part of Group III. Also other ten diabetic rats; part of Group IV received 400 mg/kg of ginger orally for six weeks. The last ten diabetic rats in Group V received oral treatment for six weeks with ginger (400 mg/kg) and diosmin (100 mg/kg). Rats were sacrificed after the six weeks, and kidney tissue and blood samples were taken for several biochemical analyses. Both an ultrastructural investigation and hemotoxylin and eosin staining were performed. The desmin and Nrf2 immunohistochemical markers were used. There were statistical and morphometric studies performed.

**Results:** the diosmin plus ginger therapy showed a considerable decrease in blood glucose, urea, and creatinine in contrast to rats with diabetes. After six weeks, there was a noticeable recovery in the levels of malondialdehyde (MDA), catalase (CAT) and superoxide dismutase (SOD). Histopathological analysis verified that group V likewise had apparently normal renal tissue architecture. When compared to the diabetic group, area% of Desmin in group V was much lower, while its area% of Nrf2 was significantly greater.

**Conclusion:** Ginger treatment alone or in combination with diosmin exhibited more pronounced improvement in diabetic nephropathy.

**Keywords:** Diosmin; Ginger; Diabetic Nephropathy; Nrf2; Desmin.

## Introduction

Diabetes mellitus (DM) is defined by a persistently high blood sugar level coupled with abnormalities in the metabolism of carbohydrates, lipids, and proteins. These abnormalities can be caused by deficiencies in either alter the release of insulin, the function of insulin, or both <sup>(1)</sup>. Elevated glucose concentrations in the cellular area cause oxidative stress because they enhance the formation of free radicals and lower natural antioxidant defenses. The development of diabetic problems is closely linked to oxidative stress, which is primarily caused by the production of free radicals that cause tissue damage. Diabetes is a major contributor to kidney failure, coronary heart disease, stroke, lower limb amputation, and blindness <sup>(2)</sup>. Various insulin formulations and oral hypoglycemic drugs, such as sulfonylureas, are being utilized to treat Type 2 diabetes. These medications provide only moderate protection against renal advancement, despite their substantial benefits in managing hyperglycemia <sup>(3)</sup>.

Ginger, or *Zingiber officinale* roscoe, is a member of the Zingiberaceae family and is a common spice in cooking. *Zingiber officinale* has major biologically active chemicals such paradols, shogaols, and gingerols, but a chemical examination reveals that it contains over 400 distinct compounds. Common illnesses including headaches, nausea, and colds are frequently treated with ginger alone <sup>(4)</sup>. *Zingiber officinale* has been shown to have a variety of therapeutic effects in mouse experiments, including anti-emetic properties in cancer treatment <sup>(5)</sup> and anti-inflammatory properties against inflammation-related disorders such colitis <sup>(6)</sup>. Many in vitro and in vivo studies have looked at ginger's and its compounds' anti-oxidative properties. The results showed that ginger decreased induced lipid peroxidation and markedly increased serum glutathione and antioxidant enzyme levels <sup>(7)</sup>.

Diosmin (diosmetin 7-O-rutinoside, C<sub>28</sub>H<sub>32</sub>O<sub>15</sub>) is a naturally present flavone glycoside that is commonly found in citrus trees that are part of the rutaceae family, which includes tangerines (*Citrus reticulata*). Its molecular weight is 608.549 g/mol. It presents significant pharmacological properties like anti-oxidant, anti-inflammatory, anti-hyperglycemic, anti-mutagenic, anti-rheumatic, and anti-allergic properties <sup>(8)</sup>. Diosmin has been shown in a recent study to protect against cadmium-

induced liver damage <sup>(9)</sup>, cerebral ischemia <sup>(10)</sup> and memory impairment brought on by traumatic brain damage <sup>(11)</sup>.

The current study aimed to: assess zingerone's protective effect against induced-diabetes; assess Diosmin's protective effect against diabetes-mediated nephropathy; assess the protective effect of their combination; and compare the renoprotective effect of both drugs with induced-diabetes through changes in the kidney's histology.

## Materials and Methods

### Animals

we used fifty adult male albino rats, ranging 200–250 g of weight that we bought from the Faculty of Medicine's animal house at Cairo University. The rats were housed in standard conditions,  $24 \pm 10^{\circ}\text{C}$ ,  $65 \pm 5\%$  humidity, and a reversible 12-hour light/dark cycle. All rats had access to clean water to drink and eat. The Benha Ethics Committee's approval was granted in compliance with the rules of ethics for experimental animals (Ethical Approval NO.: RC 16-9-2023).

### Chemicals

Alloxan monohydrate powder (Cat no. A7413- 10g) was supplied from Sigma-Aldrich Chemie GmbH.

Diosmin powder (CAS no. 520-27-4) was obtained from Sigma-Aldrich Chemie GmbH.

Ginger was obtained from MEPACO Company (Sharquia, Egypt) as 30 tablets (each tablet has 400 mg of ginger rhizome powder).

### Experimental design

Five experimental groups were randomly selected from among fifty rats; each group was included:

**Group I (control group):** consisted of ten rats. The rats were separated into five subgroups, each consisting of two rats as follows:

Subgroup Ia: rats were given a standard diet for six weeks.

Subgroup Ib: rats was injected intraperitoneally with 1 ml of normal saline as single dose equal the amount used to dissolve alloxan monohydrate.

Subgroup Ic: rats were given diosmin 100 mg/kg (dissolved in saline in total volume of 1 ml) orally through gastric gavage every day for six weeks <sup>(12)</sup>

Subgroup Id: rats were received with 400 mg/kg of ginger (dissolved in saline in total volume of 1 ml) orally every day for six weeks through gastric gavage <sup>(13)</sup>.

Subgroup Ie: rats were given diosmin 100 mg/kg (dissolved in saline in total volume of 1 ml) in addition to 400 mg/kg of ginger (dissolved in saline in total volume of 1 ml) orally by gastric gavage for six weeks.

#### **Group II (Diabetic group):**

All of the other forty rats were used to induce Diabetes. Rats, that had been starved all night, were given 150 mg/kg of alloxan monohydrate (dissolved in saline in total volume of 1 ml) through an intraperitoneal injection. The blood glucose monitoring equipment (SD biosensor Inc. Korea) was used to measure the blood glucose levels from the tail vein after a fast of 72 hours. Rats that tested for blood glucose and recorded more than 250 mg/dl were chosen to complete this study. Since alloxan induces fatal hypoglycemia six hours after injection, the rats were fed a 20% glucose solution for the next twenty-four hours in 50 ml small water feeding bottles <sup>(14)</sup>. After that, the rats with developed diabetes were fed a regular diet for a period of six weeks.

As a result, ten diabetic rats were housed for six weeks without receiving any medication and considered as diabetic group as chronic hyperglycemia is typically the precursor to diabetic nephropathy <sup>(14)</sup>. The remaining thirty diabetic rats were used in group III, IV and V.

**Group III (Dio group):** ten diabetic rats were received diosmin treatment (100 mg/kg orally every day for six weeks as in subgroup Ic) after diabetes was well established (induction of diabetes by alloxan as mentioned in group II).

**Group IV (Gin group):** ten diabetic rats were received ginger (400 mg/kg orally every day for six weeks as in subgroub Id) after diabetes was well established (induction of diabetes by alloxan as mentioned in group II).

**Group V (D+G group):** ten diabetic rats received combination of ginger and diosmin in a dose as in subgroup Ie after diabetes was well established (induction of diabetes by alloxan as mentioned in group II).

The rats were anesthetized at the end of the trial by breathing in diethyl ether. After the rats were sacrificed, blood sample (5 ml.) was collected from tail vein of each rat in the plain tube for biochemical analysis to determine blood glucose levels and renal functions. Specimens were prepared for light and electron microscopy investigations following kidney dissection.

#### ***Biochemical measurements***

The blood glucose levels were measured using the blood glucose monitoring equipment. Blood samples were centrifuged for 20 minutes after being allowed to clot for two hours at room temperature. The serum was easily removed and stored at -20 °C. The enzymatic, colorimetric approach was used to determine the urea level <sup>(15)</sup>. Quantitative reagent kit also used to test creatinine level <sup>(16)</sup>. All the kits were purchased from Biodiagnostics Co. (Cairo, Egypt).

#### **Oxidative stress and antioxidant biomarkers**

After being dissected, sliced frozen kidneys were mixed with distilled water until smooth. A clear kidney homogenate supernatant was used to identify antioxidant and oxidative stress indicators.

Considering that the quantity of malondialdehyde formed when polyunsaturated fatty acids break down, the MDA (lipid peroxidation) (MDA Elabscience Kit Catalog No. E-EL-0060, USA) was estimated <sup>(17)</sup>. Additionally, the level of superoxide dismutase activity (SOD antioxidant enzyme) (SOD Elabscience Kit Catalog no. E-EL-R1424, USA) was measured using the Marklund and Marklund approach <sup>(18)</sup> because pyrogallol inhibits auto-oxidation by catalyzing the breakdown of superoxide. Monitoring the breakdown of hydrogen peroxide enzyme, which is

catalyzed by potassium permanganate, allowed researchers to determine the degree of antioxidant enzyme of CAT (CAT Elabscience Kit Catalog No. E-BC-K031-M, USA) <sup>(19)</sup>.

### **Histological and immunohistochemical studies**

Right kidney specimens were embedded in formalin saline 10% buffer. Paraffin sections ranging in thickness from 5 to 7  $\mu$ m were arranged and placed on glass slides for staining with H and E <sup>(20)</sup>. Sections of tissue that had been deparaffinized, were treated with 3% H<sub>2</sub>O<sub>2</sub> for thirty minutes. Following this, they were incubated with anti-Desmin and anti-Nrf2 (Nuclear factor erythroid 2-related factor 2) for one hour at 37 °C as directed by the manufacturer. The method used to complete was taken from Jakkson, & Blqthe <sup>(21)</sup>.

The primary Rabbit Recombinant Monoclonal Desmin antibody was used (ab227651, Abcam [SP138], Waltham, MA, 1:100). Brown color in the cytoplasm of glomerular podocytes served as an indicator of podocyte stress <sup>(22)</sup>. Nrf2 (PA5-27882, Thermo Fisher Scientific, Rockford, USA, 1:100) is one of oxidative stress response marker that binds to antioxidant response elements in the promoter regions. The positive reaction was brown color in the cytoplasm of renal tubules <sup>(23)</sup>.

In the Anatomy and Embryology department of the Faculty of Medicine at Benha University, H&E and Immunohistochemical sections were examined and captured on camera using a light microscope (Olympus CX 41, Japan) and an attached camera (Olympus E 330, Japan).

### **Ultrastructural preparations**

Left kidney samples of approximately 1 mm<sup>3</sup> were sliced and preserved in a 3% glutaraldehyde–formaldehyde solution for ultrastructural examinations. After being cleaned with PB (pH 7.4), these samples were post-fixed for an hour at 4 °C in 1% isotonic osmium tetroxide post. The sections were processed for electron microscopic analysis. We stained the semi-thin slices with toluidine blue in order to determine the target location. After that, we used glass blades from an ultramicrotome to create ultrathin slices <sup>(24)</sup>. After staining the specimens with lead citrate and uranyl acetate, we examined the samples under an electron microscope using the JEOL

(JEM-100 SX Akishima, Tokyo, Japan) transmission. It was carried out at the Benha University Faculty of Medicine's Electron Microscopy Unit at the Histology Department.

### Statistical analysis

The % area of Desmin and Nrf2 positive expression in the innunohistochemical renal sections at 400× magnification was determined using the Image J software analyzer computer. Using SPSS v28 (IBM Inc., Armonk, NY, USA) for the statistical analysis, All information, including the mean Area (%) of positive immunohistochemistry expression levels of Nrf2 and Desmin in immune stained sections was represented as mean±SD for each group. Also, blood glucose, serum urea and creatinine levels, MDA, SOD and CAT levels were represented as mean±SD for each group .The two groups were compared using the ANOVA (F) test. The qualitative variables were analyzed using the Chi-square test, which were given as frequency and percentage (%). P-values of less than 0.05 and higher than 0.05, respectively, indicated that the results were statistically significant and nonsignificant at that point.

### Results

#### *Biochemical measurements*

Diabetic group, Dio and Gin groups had significantly increase in glucose levels than group I. Dio, Gin and D+G groups had significantly decreased in glucose levels than Diabetic group. Gin and D+G groups had significantly lower glucose levels than Dio group (table 1).

In comparison to control group, the urea level increased significantly in Dio, Gin and D+G groups. In comparison to Diabetic group, the urea level decreased significantly in in Dio, Gin and D+G groups. In comparison to Dio group, the urea level decreased significantly in in Gin and D+G groups, and it was significantly lower in D+G group when compared to Gin group (table 1).

Diabetic and Dio groups had significantly increased in creatinine levels than control group. In comparison to Diabetic group, the levels of creatinine in Dio, Gin



and D+G groups were significantly decreased. It was significantly decreased in D+G group when compared to Dio group (table 1).

### **Oxidative stress and antioxidant biomarkers**

In comparison to control group, MDA was significantly increased in Diabetic group Dio and Gin groups. MDA in Gin and D+G groups was significantly decreased than in Diabetic group. Furthermore, MDA was significantly increased in Gin group as compared to D+G group and significantly decreased in Gin and D+G groups as compared to Dio group.

In comparison to control group, the SOD levels in Diabetic, Dio, and Gin and D+G groups were significantly reduced. In comparison to Diabetic group, the SOD levels in Gin and D+G groups were significantly increased. SOD level in D+G group was significantly decreased than that of Dio and Gin groups.

In comparison to group I, the levels of catalase in Diabetic and Dio groups were significantly decreased. Gin and D+G groups were significantly increased of catalase than in Diabetic group. Catalase level of D+G group was significantly increased comparing to Dio group.

### **Hematoxylin and eosin stain**

Kidney segment examination of every subgroup in control group produced the identical histological results; hence they were all regarded as a control group. H and E-stained slices of the control group showed the renal tubules and the renal corpuscle in the renal cortex. The renal corpuscle consisted of renal glomeruli, which were surrounded by Bowman's capsule and had a tuft of capillaries. The urinary gap is what separated Bowman's capsule from the glomerulus. A single layer of basic squamous epithelium coated the parietal covering of Bowman's capsule. Cuboidal cells with acidophilic cytoplasm bordered the proximal convoluted tubules (PCT). They had centrally located, vesicular, rounded nuclei. The apical brush borders and poorly defined boundaries were seen in the cells. In the distal convoluted tubules (DCT), the lumen was broad. The cuboidal cells lined DCT and had spherical vesicular nuclei (Figure.1)

However, diabetic group had shrinkage in glomerular capillaries with wide urinary space. The majority of the PCT had dilated lumen and missing of apical brush border. Renal tubules had intraluminal debris, which indicated desquamated cells. Both the proximal and distal tubular cells showed vacuolated cytoplasm (Figure.2).

On the other hand, administration of Diosmin in group III improved glomerular structure and increased Bowman's space. Most likely, some of the renal tubules were normal. Certain tubular cells exhibited dark stained nuclei and cytoplasmic vacuolizations (Figure.3).

The diabetes histological alterations in Gin group improved; there was normal Bowman's space and about normal convoluted tubules with some vacuolated cells. Peritubular congested blood vessels were seen (Figure.4). D+G group resembled the control group in most respects with apparently normal of the glomerular capillaries and urinary space. PCT had rounded nuclei with apical brush borders. DCT lined with cubical cells with round nuclei. Congested peritubular blood vessels were observed (Figure.5).

### Ultrastructural findings

The glomerular capillaries in sections of control rats were lined with fenestrated endothelial cells with basement membrane. Glomerular basement membrane showed up with a consistent and homogenous thickness. Clear short secondary foot processes of Podocytes with renal filtration slits in between were seen around the glomerular capillaries. The glomerular capillaries' endothelial cells were intimately associated with the secondary processes to constitute the filtration barrier (Figure.6). The proximal convoluted tubule lined with a tall cuboidal shape cells with euchromatic nucleus resting on uniform basal lamina. They had narrow lumen and many apical microvilli. The sections displayed many lysosomes and mitochondria. Lateral membrane interdigitation was observed (Figure.7). Distal tubule cells were cubical in shape with a euchromatic nucleus and a few short apical microvilli. The basal part had basal infoldings and mitochondria. Intact junctional complex between the cells were seen (Figure.8).

The filtration barrier of the glomerulus in diabetic group (group II) showed vacuolated capillary endothelium. The glomerular basement membrane was heterogeneous thickened. Secondary foot processes were retracted, broadened, and

shortened as a result of the podocyte effacement with obliteration of the filtration slits (Figure.9). Missing microvilli were seen in the PCT lining cells. The nucleus had an indented appearance and seemed heterochromatic. Destructed cristae caused the mitochondria to swell. Numerous vacuoles were also seen in the rarified cytoplasm (Figure.10). In DCT cells, cytoplasm had been extruded into the tubular lumen. Its nucleus was asymmetry with the expanded chromatin. Cytoplasmic vacuoles and destructed mitochondria were seen. There was separated lateral membrane (Figure.11).

Rats treated with diosmin (Group III) showed decreased thickness of glomerular homogeneous basement membrane in their renal cortex, along with obliterated capillary endothelium fenestrae. The podocytes' secondary foot processes were detected with area of effacement (Figure.12). Proximal convoluted tubule cells had regular euchromatic nuclei. Loss of microvilli in large area was observed, with cytoplasmic vacuoles and few destructed mitochondria (Figure.13). Cuboidal cells of DCT had round nucleus with extended chromatin. The luminal border showed few apical short microvilli. Cytoplasmic vacuoles DCT showed less cytoplasmic vacuolization with normal mitochondria of various sizes and shapes (Figure.14).

Gin group exhibited more improvement, where a fenestrated capillary endothelium was observed. The glomerular basement membrane thickness was apparently normal with regularly arranged, interdigitating pedicels (Figure.15). Proximal tubule cells contained euchromatic round nucleus and few cytoplasmic vacuoles with focal loss of apical microvilli (Figure.16). Cuboidal cells of distal tubules had euchromatic nucleus. Basal folding had mitochondria in between. Short microvilli lining the lumen were seen (Figure.17).

The renal cortex ultrastructure of D+G group, the glomerular filtration barrier formed of fenestrated capillary endothelium resting on regular glomerular basement. The shape of podocytes foot processes was emerged almost to control group (Figure.18). PCT, and DCT were all preserved and resembled the ultrastructure of the control group almost exactly. It was observed PCT with regular euchromatic nucleus apical microvilli and normal mitochondria. Few cytoplasmic vacuoles were detected (Figure. 19). DCT showed cuboidal cells with regular normal nucleus, mitochondria, and characteristic short microvilli lining lumen. There was an improvement in microvilli of PCT (Figure. 20).

### Immunohistochemical studies

It was found that all subgroups of control group had minimal expressed desmin immunoexpression in the glomerular podocytes, but diabetic group had obvious positive expression of desmin. Desmin expression was less obvious in Dio and Gin groups. Area % of Desmin was significantly higher in Diabetic, Dio and Gin groups compared to control group. Area % of Desmin was significantly decreased in Dio, Gin and D+G groups compared to Diabetic group. Area % of Desmin revealed significantly decreased in D+G group compared to Dio group (table 3; Figure.21a-e).

The kidneys from each group are depicted in table (3) and Figures (22a-e), corresponding to immunohistochemical analyses of Nrf2. In renal tubular epithelial cells, results from control group revealed a less obvious expression of Nrf2. The renal tissues of Diabetic group had minimal Nrf2 positive regions. Area % of Nrf2 was significantly decreased in Diabetic and Dio groups compared to control group. Renal tubular epithelial cells of Dio, Gin and D+G groups also showed obvious Nrf2 expression. Area % of Nrf2 was significantly increased in Dio, Gin and D+G groups compared to Diabetic group. Area % of Nrf2 was significantly increased in D+G group compared to Dio and Gin groups.

### Discussion

One of the largest healthcare problems in the world is diabetes mellitus (DM), and as the disease's prevalence rises, so do its costs. For example, according to Abhinav et al. <sup>(25)</sup>, 463 million individuals globally (aged 18 to 99) were impacted by DM in 2019. The anticipated number of diabetics in Egypt was 10.9 million. By 2030, this number is expected to rise to 13 million, and by 2045, it is expected to reach 20 million <sup>(3)</sup>.

Long-term hyperglycemia in diabetes mellitus results in elevated advanced glycation end products (AGEs). AGEs build up in tissue, which leads to nephropathy, target protein and lipid denaturation, and functional deterioration <sup>(26)</sup>.

The goal of the current study was to examine the potential renoprotective effects of diosmin and ginger, either separately or in combination, against oxidative

stress caused by hyperglycemia in diabetic nephropathy caused by alloxan. Because alloxan can have cytotoxic effects on the pancreatic  $\beta$  cells, it is a commonly used medication for inducing diabetes mellitus in experimental rats <sup>(27)</sup>. The current study's alloxan-treated group exhibits the traditional diabetic nephropathy signs, namely hyperglycemia consumption. Ginger and/or diosmin therapy tend to reduce the rise in blood sugar. Our findings support the research conducted by other researchers on the potential antidiabetic benefits of natural compounds <sup>(28, 29)</sup>. Sultana et al., <sup>(30)</sup> explained that ginger can antagonize suppression of insulin release induced through serotonin receptors.

Protein metabolism's primary byproduct is urea, whereas skeletal muscle mass is represented by creatinine, a byproduct of creatinine and phosphoserine. Reduced skeletal muscle volume indicates fewer insulin target sites, which results in insulin resistance and Type 2 diabetes <sup>(31)</sup>. Generally speaking, renal impairment is indicated by elevated urea and creatinine values. In this study, diabetic rats showed considerably higher levels of kidney damage indicators, such as urea and serum creatinine, than the control group. Our findings concur with those published by Sekiou et al. <sup>(32)</sup>, who discovered that these markers are markedly increased in rats with diabetes caused by alloxan. The high level of protein breakdown may be the cause of the diabetic rats' group's notable rise in those indicators of renal damage. On the other hand, we have shown that the treatment of Diosmin or/and Ginger significantly decreased blood creatinine and urea in diabetic rats, indicating a considerable reduction in protein breakdown. Our results are in good agree with earlier research <sup>(29, 33)</sup>, which showed that rats given Diosmin or Ginger, respectively, saw a substantial drop in urea and creatinine.

In this work, rats given alloxan to induce diabetes showed elevation in MDA generation and a considerable reduce in SOD and CAT activity when compared to the control group. This is in line with previous studies <sup>(32, 34)</sup>, which showed that after being exposed to alloxan, MDA levels rose while SOD and CAT levels dropped. Intracellular auto-oxidation of glucose to glycol is brought on by AGEs. More free radicals are produced when the reactive products combine with the amino groups of extracellular and intracellular proteins to create AGEs. Lipid peroxidation and MDA formation are the results of increased oxidative stress <sup>(35)</sup>.

Conversely, In groups treated with diosmin or/and ginger, there was a considerable decrease in MDA levels accompanied by significantly higher SOD and catalase activity, suggesting that the antioxidant system has been activated to counterbalance the excess ROS generated. These results concur with those of previous research that also revealed treatment with diosmin or/ and ginger significantly increase SOD and catalase activity and significantly decrease in MDA level demonstrating its potent antioxidant action and lowers cell death by binding with free radicals <sup>(8, 36)</sup>.

Under hematoxylin and eosin staining, the renal cortex sections of the alloxan-induced diabetic group in this study had dilated Bowman space and glomerular abnormalities. Detached cells were present into the lumen tubules. The results that were reported earlier agreed with these findings <sup>(37)</sup>. Electron analyses of the renal cortex support these conclusions. In addition to podocyte effacement, the glomerular basement membrane was unevenly thickened. Abrupt loss of apical microvilli was seen in PCT cells. Cytoplasmic vacuolations, enlarged mitochondria, and pyknotic nuclei were all seen. The intracellular space between the cells exhibiting cytoplasmic vacuolations increased in DCT cells. These findings are agree with earlier research <sup>(38, 39)</sup>, who noted kidney alterations in diabetes caused by streptozotocin. This might be because excessive extracellular matrix formation and mechanical stretching are brought on by hyperglycemia's induction of hyperfiltration and hyperperfusion <sup>(40)</sup>.

In the present study, Gin group showed improvement of diabetic-induced degenerative changes after administration of ginger. Ultrastructurally, a regular thickness of basement membrane and attenuation of effacement with few vacuolization in PT were seen. Similar results were reported in previous studies <sup>(13, 41, 42)</sup>. This could be attributed to the hypoglycemic effect of ginger. Ginger was shown to reduce inflammatory markers in diabetic individuals <sup>(43)</sup>. In addition, it can improve the oxidative stress and has antiapoptotic role <sup>(13, 44)</sup>.

Rats given a combination of diosmin and ginger in our investigation showed a more noticeable improvement in diabetic-induced renal degenerative disease. This study aligns with the goal of creating a cosmeceutical preparation that carries Diosmin in conjunction with various essential oils, like Zingiber officinale (ginger oil), to function as both preventive and curative dermal novel preparations, aimed at battling

harmful environmental factors that impact the skin. The generated lipid colloidal carriers (LCCs) were evaluated, and the addition of the mentioned ginger oil was found to improve the Diosmin action in both in-vitro and in-vivo investigations <sup>(45)</sup>.

Desmin function is to raise the mechanical resistance of the cells, when its expression increases in podocytes it represents an indicator of morphological changes of these podocytes in reaction to injury <sup>(46)</sup>. Immunohistochemical staining for Desmin of diabetic group revealed, intense positive immune reaction of glomerular podocytes as compared to the control group. Moreover, moderate immune reaction of glomerular podocytes was observed in groups treated with diosmin or ginger. Administration of diosmin and ginger in diabetic rats had weak reaction similar to control group. Previous studies suggested that diosmin and ginger have a strong antioxidant, anti-inflammatory, antifibrotic, immunomodulating and antiapoptotic properties <sup>(8, 36)</sup>.

To further explain the route, the immunohistochemistry of Nrf2 was examined. In renal tubular epithelial cells, Diabetic group had weak Nrf2 positive regions. These findings were consistent with Abdul-Hamid et al. <sup>(38)</sup> and Chen et al. <sup>(47)</sup> who found that Streptozotocin-induced diabetes caused significant decreases in Nrf2 level. Renal tubular epithelial cells of groups treated with Diosmin and Ginger also showed considerable Nrf2 expression. Compared to other groups, D+G group had higher intense of positive regions for Nrf2 which showed a protective effect of diosmin and ginger against diabetic-induced nephropathy. It was shown in an animal study that Nrf2 plays a critical role in reducing the kidney damage caused by streptozotocin. This is demonstrated by the fact that Nrf2-/- mice produce more ROS than Nrf2+/- mice, and they also have more kidney damage and oxidative DNA damage <sup>(48)</sup>. Another study found that the diabetic group's spleen had much less Nrf2 expression than the control group's because hyperglycemia, which causes diabetes, stimulates Keap1 (Kelch like associated protein-1) malfunction. Another study found that the diabetic group's spleen had much less Nrf2 expression than the control group's because hyperglycemia, which causes diabetes, stimulates Keap1 (Kelch like associated protein-1) malfunction <sup>(49)</sup>.

## Conclusions